



Citation for published version:

Kyprianou, A, Pardo Millan, J-C & Watson, A 2014, 'The extended hypergeometric class of Lévy processes', *Journal of Applied Probability*, vol. 51A, pp. 391 - 408. <https://doi.org/10.1239/jap/1417528488>

DOI:

[10.1239/jap/1417528488](https://doi.org/10.1239/jap/1417528488)

Publication date:

2014

Document Version

Peer reviewed version

[Link to publication](#)

(C) 2014 Applied Probability Trust. The published version is available via:
<http://dx.doi.org/10.1239/jap/1417528488>

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

The extended hypergeometric class of Lévy processes

A. E. Kyprianou^{*} J. C. Pardo[†] A. R. Watson[‡]

23rd September 2014

With a view to computing fluctuation identities related to stable processes, we review and extend the class of hypergeometric Lévy processes explored in Kuznetsov and Pardo [17]. We give the Wiener–Hopf factorisation of a process in the extended class, and characterise its exponential functional. Finally, we give three concrete examples arising from transformations of stable processes.

1 Introduction

The simple definition of a Lévy process—a stochastic process with stationary independent increments—has been sufficient to fuel a vast field of study for many decades, and Lévy processes have been employed in many successful applied models. However, historically there have been few classes of processes for which many functionals could be computed explicitly. In recent years, the field has seen a proliferation of examples which have proved to be more analytically tractable; in particular, we single out spectrally negative Lévy processes [20], Lamperti-stable processes [6, 9], β - and θ -processes [15, 16], and finally the inspiration for this work, hypergeometric Lévy processes [17, 18, 25]. It is also worth mentioning that the close relationship which appears to hold between hypergeometric Lévy processes and stable processes has also allowed the computation of several identities for the latter; see [17, 25, 26].

^{*}University of Bath, UK. a.kyprianou@bath.ac.uk

[†]CIMAT, Mexico. jcpardo@cimat.mx

[‡]University of Zürich, Switzerland. alexander.watson@math.uzh.ch. Parts of this work were completed while the third author was at the University of Bath, UK, and at CIMAT, Mexico.

Keywords and phrases: Lévy processes, hypergeometric Lévy processes, extended hypergeometric Lévy processes, Wiener–Hopf factorisation, exponential functionals, stable processes, path-censored stable process, conditioned stable processes, hitting distributions, hitting probabilities.

MSC 2000 classification: 60G51, 60G18, 60G52.

In this work, we review the hypergeometric class of Lévy processes introduced by Kuznetsov and Pardo [17], and introduce a new class of extended hypergeometric processes which have many similar properties. In particular, for an extended hypergeometric process ξ we compute the Wiener–Hopf factors and find that its ladder height processes are related to Lamperti-stable subordinators; and we are able to characterise explicitly the distribution of the exponential functional of ξ/δ for any $\delta > 0$. We also give three examples of processes connected via the Lamperti representation to α -stable processes, which fall into the hypergeometric class when $\alpha \leq 1$, and into the extended hypergeometric class when $\alpha > 1$, and give some new identities for the stable process when $\alpha > 1$.

We will first discuss the results of Kuznetsov and Pardo [17]. For a choice of parameters $(\beta, \gamma, \hat{\beta}, \hat{\gamma})$ from the set

$$\mathcal{A}_{\text{HG}} = \{\beta \leq 1, \gamma \in (0, 1), \hat{\beta} \geq 0, \hat{\gamma} \in (0, 1)\},$$

we define

$$\psi(z) = -\frac{\Gamma(1 - \beta + \gamma - z)}{\Gamma(1 - \beta - z)} \frac{\Gamma(\hat{\beta} + \hat{\gamma} + z)}{\Gamma(\hat{\beta} + z)},$$

which we view as a meromorphic function on \mathbb{C} . We say that a Lévy process ξ is a member of the *hypergeometric class* of Lévy processes if it has Laplace exponent ψ , in the sense that

$$\mathbb{E}[e^{z\xi_1}] = e^{\psi(z)}, \quad z \in i\mathbb{R}. \quad (1)$$

Note that, in general, when the Laplace exponent ψ of a Lévy process ξ is a meromorphic function, the relation (1) actually holds on any neighbourhood of $0 \in \mathbb{C}$ which does not contain a pole of ψ ; thus, in this article we will generally not specify the domain of Laplace exponents which may arise.

In [17], it is shown that for any choice of parameters in \mathcal{A}_{HG} , there is a Lévy process with Laplace exponent ψ , and they find its Wiener–Hopf factorisation, in the following sense.

The (*spatial*) *Wiener–Hopf factorisation* of a Lévy process ξ with Laplace exponent ψ consists of the equation

$$\psi(z) = -\kappa(-z)\hat{\kappa}(z), \quad z \in i\mathbb{R},$$

where κ and $\hat{\kappa}$ are the Laplace exponents of subordinators H and \hat{H} , respectively, this time in the sense that $\mathbb{E}[e^{-\lambda H_1}] = e^{-\kappa(\lambda)}$ for $\text{Re } \lambda \geq 0$. The subordinators H and \hat{H} are known as the ascending and descending ladder heights, and are related via a time-change to the running maximum and running minimum of the process ξ . For more details, we refer the reader to [22, Chapter 6]. The insight into the structure of ξ given by the Wiener–Hopf factorisation allows one to simplify first passage problems for ξ ; see [22, Chapter 7] for a collection of results.

The paper [17] computes that

$$\kappa(z) = \frac{\Gamma(1 - \beta + \gamma + z)}{\Gamma(1 - \beta + z)}, \quad \hat{\kappa}(z) = \frac{\Gamma(\hat{\beta} + \hat{\gamma} + z)}{\Gamma(\hat{\beta} + z)},$$

thus demonstrating that the ascending and descending ladder height processes are Lamperti-stable subordinators (see [6]).

Kuznetsov and Pardo also consider the exponential functional of a hypergeometric Lévy process ξ . For each $\delta > 0$, the random variable

$$I(\xi/\delta) = \int_0^\infty e^{-\xi_t/\delta} dt$$

is a.s. finite provided that ξ drifts to $+\infty$. This random variable is known as the *exponential functional* of the Lévy process ξ , and it has been studied extensively in general; the paper of Bertoin and Yor [3] gives a survey of the literature, and mentions, among other aspects, applications to diffusions in random environments, mathematical finance and fragmentation theory. In the context of self-similar Markov processes, the exponential functional appears in the entrance law of a pssMp started at zero (see, for example, Bertoin and Yor [2]), and Pardo [31] relates the exponential functional of a Lévy process to envelopes of its associated pssMp; furthermore, it is related to the hitting time of points for pssMps, and we shall make use of it in this capacity in our example of subsection 4.2.

For the purpose of characterising the distribution of $I(\xi/\delta)$, its Mellin transform

$$\mathcal{M}(s) = \mathbb{E}[I(\xi/\delta)^{s-1}],$$

is useful. For ξ in the hypergeometric class, \mathcal{M} was calculated by [17] in terms of gamma and double gamma functions; we will recall and extend this section 3.

We now give a brief outline of the main body of the paper. In section 2, we demonstrate that the parameter set \mathcal{A}_{HG} may be extended by changing the domains of the two parameters β and $\hat{\beta}$, and find the Wiener–Hopf factorisation of a process ξ in this new class, identifying explicitly the ladder height processes. In section 3, we find an expression for the Mellin transform \mathcal{M} in this new case, making use of an auxiliary hypergeometric Lévy process. In section 4, we cover three examples where the extended hypergeometric class is of use, on the way extending the result of [7] on the Wiener–Hopf factorisation of the Lamperti representation associated with the radial part of a stable process.

2 The extended hypergeometric class

We begin by defining the set of admissible parameters

$$\mathcal{A}_{\text{EHG}} = \{\beta \in [1, 2], \gamma, \hat{\gamma} \in (0, 1), \hat{\beta} \in [-1, 0]; 1 - \beta + \hat{\beta} + \gamma \geq 0, 1 - \beta + \hat{\beta} + \hat{\gamma} \geq 0\}.$$

We are interested in proving the existence of, and investigating the properties of, a Lévy process ξ whose Laplace exponent is given by the meromorphic function

$$\psi(z) = -\frac{\Gamma(1 - \beta + \gamma - z)}{\Gamma(1 - \beta - z)} \frac{\Gamma(\hat{\beta} + \hat{\gamma} + z)}{\Gamma(\hat{\beta} + z)}, \quad z \in \mathbb{C},$$

when $(\beta, \gamma, \hat{\beta}, \hat{\gamma}) \in \mathcal{A}_{\text{EHG}}$.

To allow for more concise expressions below, we also define

$$\eta = 1 - \beta + \gamma + \hat{\beta} + \hat{\gamma}.$$

We now give our main result on the existence and properties of ξ .

Proposition 1. *There exists a Lévy process ξ such that $\mathbb{E}[e^{z\xi_1}] = e^{\psi(z)}$. Its Wiener-Hopf factorisation may be expressed as*

$$\psi(z) = -(-\hat{\beta} - z) \frac{\Gamma(1 - \beta + \gamma - z)}{\Gamma(2 - \beta - z)} \times (\beta - 1 + z) \frac{\Gamma(\hat{\beta} + \hat{\gamma} + z)}{\Gamma(1 + \hat{\beta} + z)}.$$

Its Lévy measure possesses the density

$$\pi(x) = \begin{cases} -\frac{\Gamma(\eta)}{\Gamma(\eta - \hat{\gamma})\Gamma(-\gamma)} e^{-(1-\beta+\gamma)x} {}_2F_1(1 + \gamma, \eta; \eta - \hat{\gamma}; e^{-x}), & x > 0, \\ -\frac{\Gamma(\eta)}{\Gamma(\eta - \gamma)\Gamma(-\hat{\gamma})} e^{(\hat{\beta}+\hat{\gamma})x} {}_2F_1(1 + \hat{\gamma}, \eta; \eta - \gamma; e^x), & x < 0, \end{cases} \quad (2)$$

where ${}_2F_1$ is the Gauss hypergeometric function.

If $\beta \in (1, 2)$ and $\hat{\beta} \in (-1, 0)$, the process ξ is killed at rate

$$q = \frac{\Gamma(1 - \beta + \gamma)}{\Gamma(1 - \beta)} \frac{\Gamma(\hat{\beta} + \hat{\gamma})}{\Gamma(\hat{\beta})}.$$

Otherwise, the process has infinite lifetime and:

- (i) ξ drifts to $+\infty$ if $\beta > 1$, $\hat{\beta} = 0$.
- (ii) ξ drifts to $-\infty$ if $\beta = 1$, $\hat{\beta} < 0$.
- (iii) ξ oscillates if $\beta = 1$, $\hat{\beta} = 0$. In this case, ξ is a hypergeometric Lévy process.

Furthermore, the process ξ has no Gaussian component, and is of bounded variation with zero drift when $\gamma + \hat{\gamma} < 1$ and of unbounded variation when $\gamma + \hat{\gamma} \geq 1$.

Proof. We remark that there is nothing to do in case (iii) since such processes are analysed in [17]; however, the proof we give below also carries through in this case.

We will first identify the proposed ascending and descending ladder processes. Once we have shown that ψ really is the Laplace exponent of a Lévy process, this will be the proof of the Wiener-Hopf factorisation.

Before we begin, we must review the definitions of special subordinators and the \mathcal{T} -transformations of subordinators. Suppose that v is the Laplace exponent of a subordinator H , in the sense that $\mathbb{E}[e^{-zH_1}] = e^{-v(z)}$. H is said to be a *special subordinator*, and v a *special Bernstein function*, if the function

$$v^*(z) = z/v(z), \quad z \geq 0,$$

is also the Laplace exponent of a subordinator. The function v^* is said to be *conjugate* to v . Special Bernstein functions play an important role in potential theory; see, for example, [34] for more details.

Again taking v to be the Laplace exponent of a subordinator, not necessarily special, we define, for $c \geq 0$, the transformation

$$\mathcal{T}_c v(z) = \frac{z}{z+c} v(z+c), \quad z \geq 0.$$

It is then known (see [13, 23]) that $\mathcal{T}_c v$ is the Laplace exponent of a subordinator. Furthermore, if v is in fact a special Bernstein function, then $\mathcal{T}_c v$ is also a special Bernstein function.

We are now in a position to identify the ladder height processes in the Wiener–Hopf factorisation of ξ . Let the (proposed) ascending factor be given, for $z \geq 0$, by

$$\kappa(z) = (-\hat{\beta} + z) \frac{\Gamma(1 - \beta + \gamma + z)}{\Gamma(2 - \beta + z)}.$$

Then some simple algebraic manipulation shows that

$$\kappa(z) = (\mathcal{T}_{-\hat{\beta}} v)^*(z),$$

with

$$v(z) = \frac{\Gamma(2 - \beta + \hat{\beta} + z)}{\Gamma(1 - \beta + \hat{\beta} + \gamma + z)},$$

provided that v is a special Bernstein function. This follows immediately from Example 2 of Kyprianou and Rivero [24], under the constraint $1 - \beta + \hat{\beta} + \gamma \geq 0$ which is included in the parameter set \mathcal{A}_{EHG} . Furthermore, we note that v is in fact the Laplace exponent of a Lamperti-stable subordinator (see [6]), although we will not use this fact.

Proceeding similarly for the descending factor, we obtain

$$\hat{\kappa}(z) = (\beta - 1 + z) \frac{\Gamma(\hat{\beta} + \hat{\gamma} + z)}{\Gamma(1 + \hat{\beta} + z)} = (\mathcal{T}_{\beta-1} \hat{v})^*(z), \quad z \geq 0,$$

where

$$\hat{v}(z) = \frac{\Gamma(2 - \beta + \hat{\beta} + z)}{\Gamma(1 - \beta + \hat{\beta} + \hat{\gamma} + z)}, \quad z \geq 0,$$

and again the function \hat{v} is a special Bernstein function provided that $1 - \beta + \hat{\beta} + \hat{\gamma} \geq 0$. As before, \hat{v} is the Laplace exponent of a Lamperti-stable subordinator.

We have now shown that both κ and $\hat{\kappa}$ are Laplace exponents of subordinators; we wish to show that the function

$$\psi(z) = -\kappa(-z)\hat{\kappa}(z)$$

is the Laplace exponent of a Lévy process. For this purpose we apply the theory of *philanthropy* developed by Vigon [35, chapter 7]. This states, in part, that it is sufficient

for both of the subordinators corresponding to κ and $\hat{\kappa}$ to be ‘philanthropists’, which means that their Lévy measures possess decreasing densities.

We recall our discussion of \mathcal{T} -transforms and special Bernstein functions. We have already stated that when v is a special Bernstein function, then so is $\mathcal{T}_c v$; furthermore, one may show that its conjugate satisfies

$$(\mathcal{T}_c v)^*(z) = \mathcal{E}_c v^*(z) + v^*(c), \quad z \geq 0,$$

where \mathcal{E}_c is the *Esscher transform*, given by

$$\mathcal{E}_c v^*(z) = v^*(z + c) - v^*(c), \quad z \geq 0.$$

The Esscher transform of the Laplace exponent of any subordinator is again the Laplace exponent of a subordinator; and if the subordinator corresponding to v^* possesses a Lévy density π_{v^*} , then the Lévy density of $\mathcal{E}_c v^*$ is given by $x \mapsto e^{-cx} \pi_{v^*}(x)$, for $x > 0$.

Returning to our Wiener-Hopf factors, we have

$$\kappa(z) = (\mathcal{T}_{-\hat{\beta}} v)^*(z) = \mathcal{E}_{-\hat{\beta}} v^*(z) + v^*(-\hat{\beta}), \quad z \geq 0,$$

where v^* is the Laplace exponent conjugate to v . Now, v is precisely the type of special Bernstein function considered in [24, Example 2]. In that work, the authors even establish that the subordinator corresponding to v^* has a decreasing Lévy density π_{v^*} . Finally, the Lévy density of the subordinator corresponding to κ is $x \mapsto e^{\hat{\beta}x} \pi_{v^*}(x)$, and this is then clearly also decreasing.

Hence, we have shown that the subordinator whose Laplace exponent is κ is a philanthropist. By a very similar argument, the subordinator corresponding to $\hat{\kappa}$ is also a philanthropist. As we have stated, the theory developed by Vigon now shows that the function ψ really is the Laplace exponent of a Lévy process ξ , with the Wiener-Hopf factorisation claimed.

We now proceed to calculate the Lévy measure of ξ . A fairly simple way to do this is to make use of the theory of ‘meromorphic Lévy processes’, as developed in Kuznetsov et al. [19]. We first show that ξ is in the meromorphic class. Initially suppose that

$$1 - \beta + \hat{\beta} + \gamma > 0, \quad 1 - \beta + \hat{\beta} + \hat{\gamma} > 0; \quad (3)$$

we will relax this assumption later. Looking at the expression for ψ , we see that it has zeroes $(\zeta_n)_{n \geq 1}$, $(-\hat{\zeta}_n)_{n \geq 1}$ and (simple) poles $(\rho_n)_{n \geq 1}$, $(-\hat{\rho}_n)_{n \geq 1}$ given as follows:

$$\begin{aligned} \zeta_1 &= -\hat{\beta}, & \zeta_n &= n - \beta, & n &\geq 2, \\ \hat{\zeta}_1 &= \beta - 1, & \rho_n &= n - \beta + \gamma, & n &\geq 1, \\ & & \hat{\zeta}_n &= \hat{\beta} + n - 1, & n &\geq 2, \\ & & \hat{\rho}_n &= \hat{\beta} + \hat{\gamma} + n - 1, & n &\geq 1, \end{aligned}$$

and that they satisfy the interlacing condition

$$\cdots < -\hat{\rho}_2 < -\hat{\zeta}_2 < -\hat{\rho}_1 < -\hat{\zeta}_1 < 0 < \zeta_1 < \rho_1 < \zeta_2 < \rho_2 < \cdots.$$

To show that ξ belongs to the meromorphic class, one applies [19, Theorem 1(v)] when ξ is killed, and [19, Corollary 2] in the unkilld case. The proof is a routine calculation using the Weierstrass representation [14, 8.322] to expand κ and $\hat{\kappa}$ as infinite products, and we omit it for the sake of brevity.

We now calculate the Lévy density. For a process in the meromorphic class, it is known that the Lévy measure has a density of the form

$$\pi(x) = \mathbb{1}_{\{x>0\}} \sum_{n \geq 1} a_n \rho_n e^{-\rho_n x} + \mathbb{1}_{\{x<0\}} \sum_{n \geq 1} \hat{a}_n \hat{\rho}_n e^{\hat{\rho}_n x}, \quad (4)$$

for some coefficients $(a_n)_{n \geq 1}$, $(\hat{a}_n)_{n \geq 1}$, where the ρ_n and $\hat{\rho}_n$ are as above. Furthermore, from [19, equation (8)], we see that

$$a_n \rho_n = -\text{Res}(\psi(z) : z = \rho_n),$$

and correspondingly for $\hat{a}_n \hat{\rho}_n$. (This remark is made on p. 1111 of [19].) From here it is simple to compute

$$a_n \rho_n = -\frac{(-1)^{n-1}}{(n-1)!} \frac{1}{\Gamma(1-\gamma-n)} \frac{\Gamma(\eta+n-1)}{\Gamma(\eta-\hat{\gamma}+n-1)}, \quad n \geq 1,$$

and similarly for $\hat{a}_n \hat{\rho}_n$. The expression (2) follows by substituting in (4) and using the series definition of the hypergeometric function.

Thus far we have been working under the assumption that (3) holds. Suppose now that this fails and we have, say, $1-\beta+\hat{\beta}+\hat{\gamma}=0$. Then $\zeta_1 = \rho_1$, which is to say the first zero-pole pair to the right of the origin is removed. It is clear that ξ still falls into the meromorphic class, and indeed, our expression for π remains valid: although the initial pole ρ_1 no longer exists, the corresponding coefficient $a_1 \rho_1$ vanishes as well. Similarly, we may allow $1-\beta+\hat{\beta}+\gamma=0$, in which case the zero-pole pair to the left of the origin is removed; or we may allow both expressions to be zero, in which case both pairs are removed. The proof carries through in all cases.

The claim about the large time behaviour of ξ follows from the Wiener-Hopf factorisation: $\kappa(0) = 0$ if and only if the range of ξ is a.s. unbounded above, and $\hat{\kappa}(0) = 0$ if and only if the range of ξ is a.s. unbounded below; so we need only examine the values of $\kappa(0)$, $\hat{\kappa}(0)$ in each of the four parameter regimes.

Finally, we prove the claims about the Gaussian component and variation of ξ . This proof proceeds along the same lines as that in [17]. Firstly, we observe using [14, formula 8.328.1] that

$$\psi(i\theta) = O(|\theta|^{\gamma+\hat{\gamma}}), \quad |\theta| \rightarrow \infty. \quad (5)$$

Applying [1, Proposition I.2(i)] shows that ξ has no Gaussian component. Then, using [14, formulas 9.131.1 and 9.122.2], one sees that

$$\pi(x) = O(|x|^{-(1+\gamma+\hat{\gamma})}), \quad x \rightarrow 0,$$

and together with the necessary and sufficient condition $\int_{\mathbb{R}} (1 \wedge |x|) \pi(x) dx < \infty$ for bounded variation, this proves the claim about the variation. In the bounded variation case, applying [1, Proposition I.2(ii)] with (5) shows that ξ has zero drift. \square

We propose to call this the *extended hypergeometric class* of Lévy processes.

Remark 2. If ξ is a process in the extended hypergeometric class, with parameters $(\beta, \gamma, \hat{\beta}, \hat{\gamma})$, then the dual process $-\xi$ also lies in this class, and has parameters $(1 - \hat{\beta}, \hat{\gamma}, 1 - \beta, \gamma)$.

Remark 3. We remark here that one may instead extend the parameter range \mathcal{A}_{HG} by moving only β , or only $\hat{\beta}$. To be precise, both

$$\mathcal{A}_{\text{EHG}}^{\beta} = \{\beta \in [1, 2], \gamma, \hat{\gamma} \in (0, 1), \hat{\beta} \geq 0; 1 - \beta + \hat{\beta} + \gamma \leq 0, 1 - \beta + \hat{\beta} + \hat{\gamma} \geq 0\}$$

and

$$\mathcal{A}_{\text{EHG}}^{\hat{\beta}} = \{\beta \leq 1, \gamma, \hat{\gamma} \in (0, 1), \hat{\beta} \in [-1, 0]; 1 - \beta + \hat{\beta} + \gamma \geq 0, 1 - \beta + \hat{\beta} + \hat{\gamma} \leq 0\}$$

are suitable parameter regimes, and one may develop a similar theory for such processes; for instance, for parameters in $\mathcal{A}_{\text{EHG}}^{\beta}$, one has the Wiener–Hopf factors

$$\kappa(z) = \frac{\Gamma(2 - \beta + \gamma + z)}{\Gamma(2 - \beta + z)}, \quad \hat{\kappa}(z) = \frac{\beta - 1 + z}{\beta - 1 - \gamma + z} \frac{\Gamma(\hat{\beta} + \hat{\gamma} + z)}{\Gamma(\hat{\beta} + z)}.$$

However, we are not aware of any examples of processes in these classes.

3 The exponential functional

Suppose that ξ is a Lévy process in the extended hypergeometric class with $\beta > 1$, which is to say either ξ is killed or it drifts to $+\infty$.

We are then interested in the *exponential functional* of the process, given by

$$I(\xi/\delta) = \int_0^\infty e^{-\xi_t/\delta} dt,$$

for $\delta > 0$. (Since ξ/δ is not in the extended hypergeometric class, we are studying exponential functionals of a slightly larger collection of processes.) This is an a.s. finite random variable under the conditions we have just outlined.

It will emerge that the best way to characterise the distribution of $I(\xi/\delta)$ is via its *Mellin transform*,

$$\mathcal{M}(s) = \mathbb{E}[I(\xi/\delta)^{s-1}],$$

whose domain of definition will be a vertical strip in the complex plane to be determined.

In the case of a hypergeometric Lévy process with $\hat{\beta} > 0$, it was shown in [17] that the Mellin transform of the exponential functional is given by

$$\mathcal{M}_{\text{HG}}(s) = C\Gamma(s) \frac{G((1 - \beta)\delta + s; \delta)}{G((1 - \beta + \gamma)\delta + s; \delta)} \frac{G((\hat{\beta} + \hat{\gamma})\delta + 1 - s; \delta)}{G(\hat{\beta}\delta + 1 - s; \delta)},$$

holds for $\text{Re } s \in (0, 1 + \hat{\beta}\delta)$, where C is a normalising constant such that $\mathcal{M}_{\text{HG}}(1) = 1$, and G is the double gamma function; see [17] for a definition of this special function.

Our goal in this section is the following result, which characterises the law of the exponential functional for the extended hypergeometric class.

Proposition 4. *Suppose that ξ is a Lévy process in the extended hypergeometric class with $\beta > 1$. Define $\theta = \delta(\beta - 1)$.*

Then, the Mellin transform \mathcal{M} of $I(\xi/\delta)$ is given by

$$\mathcal{M}(s) = c\widetilde{\mathcal{M}}(s) \frac{\Gamma(\delta(1 - \beta + \gamma) + s)}{\Gamma(-\delta\hat{\beta} + s)} \frac{\Gamma(\delta(\beta - 1) + 1 - s)}{\Gamma(\delta(\hat{\beta} + \hat{\gamma}) + 1 - s)}, \quad \operatorname{Re} s \in (0, 1 + \theta), \quad (6)$$

where $\widetilde{\mathcal{M}}$ is the Mellin transform of $I(\zeta/\delta)$, and ζ is an auxiliary Lévy process in the hypergeometric class, with parameters $(\beta - 1, \gamma, \hat{\beta} + 1, \hat{\gamma})$. The constant c is such that $\mathcal{M}(1) = 1$.

Proof. The process ξ/δ has Laplace exponent ψ_δ given by $\psi_\delta(z) = \psi(z/\delta)$. The relationship with ζ arises from the following calculation:

$$\begin{aligned} \psi_\delta(z) &= \frac{-\hat{\beta} - z/\delta}{1 - \beta + \gamma - z/\delta} \frac{\beta - 1 + z/\delta}{\hat{\beta} + \hat{\gamma} + z/\delta} \frac{\Gamma(2 - \beta + \gamma - z/\delta)}{\Gamma(2 - \beta - z/\delta)} \frac{\Gamma(1 + \hat{\beta} + \hat{\gamma} + z/\delta)}{\Gamma(1 + \hat{\beta} + z/\delta)} \\ &= \frac{-\hat{\beta} - z/\delta}{1 - \beta + \gamma - z/\delta} \frac{\beta - 1 + z/\delta}{\hat{\beta} + \hat{\gamma} + z/\delta} \tilde{\psi}_\delta(z), \end{aligned} \quad (7)$$

where $\tilde{\psi}_\delta$ is the Laplace exponent of a Lévy process ζ/δ , with ζ as in the statement of the theorem.

Denote by $f(s)$ the right-hand side of (6). The proof now proceeds via the ‘verification result’ [17, Proposition 2].

Recall that a Lévy process with Laplace exponent ϕ is said to satisfy the *Cramér condition with Cramér number θ* if there exists $z_0 < 0$ and $\theta \in (0, -z_0)$ such that $\phi(z)$ is defined for all $z \in (z_0, 0)$ and $\phi(-\theta) = 0$.

Inspecting the Laplace exponent ψ_δ reveals that ξ/δ satisfies the Cramér condition with Cramér number $\theta = \delta(\beta - 1)$.

Furthermore, ζ/δ satisfies the Cramér condition with Cramér number $\tilde{\theta} = \delta(\hat{\beta} + 1)$. It follows from [32, Lemma 2] that $\widetilde{\mathcal{M}}(s)$ is finite in the strip $\operatorname{Re} s \in (0, 1 + \tilde{\theta})$; and by the properties of Mellin transforms of positive random variables, it is analytic and zero-free in its domain of definition. The constraints in the parameter set \mathcal{A}_{EHG} ensure that $\tilde{\theta} \geq \theta$; this, together with inspecting the right-hand side of (6) and comparing again with the conditions in \mathcal{A}_{EHG} , demonstrates that $\mathcal{M}(s)$ is analytic and zero-free in the strip $\operatorname{Re} s \in (0, 1 + \theta)$.

We must then check the functional equation $f(s + 1) = -sf(s)/\psi_\delta(-s)$, for $s \in (0, \theta)$.

Apply (7) to write

$$\begin{aligned}
-\frac{s}{\psi_\delta(-s)} &= -\frac{s}{\tilde{\psi}_\delta(-s)} \frac{1-\beta+\gamma+s/\delta}{-\hat{\beta}+s/\delta} \frac{\hat{\beta}+\hat{\gamma}-s/\delta}{\beta-1-s/\delta} \\
&= \frac{\tilde{\mathcal{M}}(s+1)}{\tilde{\mathcal{M}}(s)} \frac{\delta(1-\beta+\gamma)+s}{-\delta\hat{\beta}+s} \frac{\delta(\hat{\beta}+\hat{\gamma})-s}{\delta(\beta-1)-s} \\
&= \frac{\tilde{\mathcal{M}}(s+1)}{\tilde{\mathcal{M}}(s)} \frac{\Gamma(-\delta\hat{\beta}+s)}{\Gamma(-\delta\hat{\beta}+s+1)} \frac{\Gamma(\delta(1-\beta+\gamma)+s+1)}{\Gamma(\delta(1-\beta+\gamma)+s)} \\
&\quad \times \frac{\Gamma(\delta(\hat{\beta}+\hat{\gamma})+1-s)}{\Gamma(\delta(\hat{\beta}+\hat{\gamma})-s)} \frac{\Gamma(\delta(\beta-1)-s)}{\Gamma(\delta(\beta-1)+1-s)},
\end{aligned}$$

making use of the same functional equation for the Mellin transform $\tilde{\mathcal{M}}$. It is then clear that the right-hand side is equal to $f(s+1)/f(s)$.

Finally, it remains to check that $|f(s)|^{-1} = o(\exp(2\pi|\operatorname{Im}(s)|))$, as $|\operatorname{Im} s| \rightarrow \infty$, uniformly in $\operatorname{Re} s \in (0, 1 + \theta)$. The following asymptotic relation may be derived from Stirling's asymptotic formula for the gamma function:

$$\log \Gamma(z) = z \log z - z + O(\log z), \quad (8)$$

and since Stirling's asymptotic formula is uniform in $|\arg(z)| < \pi - \omega$ for any choice of $\omega > 0$, it follows that (8) holds uniformly in the strip $\operatorname{Re} s \in (0, 1 + \theta)$; see [29, Chapter 8, §4]. We thus obtain

$$\log \left| \frac{\Gamma(\delta(1-\beta+\gamma)+s)}{\Gamma(-\delta\hat{\beta}+s)} \frac{\Gamma(\delta(\beta-1)+1-s)}{\Gamma(\delta(\hat{\beta}+\hat{\gamma})+1-s)} \right|^{-1} = O(\log s) = o(\operatorname{Im} s),$$

and comparing this with the proof of [17, Theorem 2], where the asymptotic behaviour of $\tilde{\mathcal{M}}(s)$ is given, we see that this is sufficient for our purposes.

Hence, $\mathcal{M}(s) = f(s)$ when $\operatorname{Re} s \in (0, 1 + \theta)$. \square

This Mellin transform may be inverted to give an expression for the density of $I(\xi/\delta)$ in terms of series whose terms are defined iteratively, but we do not pursue this here. For details of this approach, see [17, §4].

4 Three examples

It is well-known that hypergeometric Lévy processes appear as the Lamperti transforms of stable processes killed passing below zero, conditioned to stay positive and conditioned to hit zero continuously; see [17, Theorem 1]. In this section we briefly present three additional examples in which the extended hypergeometric class comes into play. The examples may all be obtained in the same way: we begin with a stable process, modify its path in some way to obtain a positive, self-similar Markov process, and then apply

the Lamperti transform to obtain a new Lévy process. We therefore start with a short description of these concepts.

We work with the (strictly) stable process X with scaling parameter α and positivity parameter ρ , which is defined as follows. For (α, ρ) in the set

$$\mathcal{A}_{\text{st}} = \{(\alpha, \rho) : \alpha \in (0, 1), \rho \in (0, 1)\} \cup \{(\alpha, \rho) = (1, 1/2)\} \\ \cup \{(\alpha, \rho) : \alpha \in (1, 2), \rho \in (1 - 1/\alpha, 1/\alpha)\},$$

and let X , with probability laws $(P_x)_{x \in \mathbb{R}}$, be the Lévy process with characteristic exponent

$$\Psi(\theta) = \begin{cases} c|\theta|^\alpha(1 - i\beta \tan \frac{\pi\alpha}{2} \operatorname{sgn} \theta) & \alpha \in (0, 2) \setminus \{1\}, \\ c|\theta| & \alpha = 1, \end{cases} \quad \theta \in \mathbb{R},$$

where $c = \cos(\pi\alpha(\rho - 1/2))$ and $\beta = \tan(\pi\alpha(\rho - 1/2))/\tan(\pi\alpha/2)$; by this we mean that $E_0[e^{i\theta X_1}] = e^{-\Psi(\theta)}$. This Lévy process has absolutely continuous Lévy measure with density

$$c_+ x^{-(\alpha+1)} \mathbb{1}_{\{x>0\}} + c_- |x|^{-(\alpha+1)} \mathbb{1}_{\{x<0\}}, \quad x \in \mathbb{R},$$

where

$$c_+ = \frac{\Gamma(\alpha + 1)}{\Gamma(\alpha\rho)\Gamma(1 - \alpha\rho)}, \quad c_- = \frac{\Gamma(\alpha + 1)}{\Gamma(\alpha\hat{\rho})\Gamma(1 - \alpha\hat{\rho})}$$

and $\hat{\rho} = 1 - \rho$.

The parameter set \mathcal{A}_{st} and the characteristic exponent Ψ represent, up a multiplicative constant in Ψ , all (strictly) stable processes which jump in both directions, except for Brownian motion and the symmetric Cauchy processes with non-zero drift.

The choice of α and ρ as parameters is explained as follows. X satisfies the α -scaling property, that

$$\text{under } P_x, \text{ the law of } (cX_{tc^{-\alpha}})_{t \geq 0} \text{ is } P_{cx}, \quad (9)$$

for all $x \in \mathbb{R}$, $c > 0$. The second parameter satisfies $\rho = P_0(X_t > 0)$.

A *positive self-similar Markov process (pssMp)* with *self-similarity index* $\alpha > 0$ is a standard Markov process $Y = (Y_t)_{t \geq 0}$ with filtration $(\mathcal{G}_t)_{t \geq 0}$ and probability laws $(P_x)_{x>0}$, on $[0, \infty)$, which has 0 as an absorbing state and which satisfies the scaling property (9) (with Y in place of X). Here, we mean “standard” in the sense of [4], which is to say, $(\mathcal{G}_t)_{t \geq 0}$ is a complete, right-continuous filtration, and Y has càdlàg paths and is strong Markov and quasi-left-continuous.

In the seminal paper [27], Lamperti describes a one to one correspondence between pssMps and Lévy processes, which we now outline. It may be worth noting that we have presented a slightly different definition of pssMp from Lamperti; for the connection, see [36, §0].

Let $S(t) = \int_0^t (Y_u)^{-\alpha} du$. This process is continuous and strictly increasing until Y reaches zero. Let $(T(s))_{s \geq 0}$ be its inverse, and define

$$\xi_s = \log Y_{T(s)} \quad s \geq 0.$$

Then $\xi := (\xi_s)_{s \geq 0}$ is a Lévy process started at $\log x$, possibly killed at an independent exponential time; the law of the Lévy process and the rate of killing do not depend on the value of x . The real-valued process ξ with probability laws $(\mathbb{P}_y)_{y \in \mathbb{R}}$ is called the *Lévy process associated to Y* , or the *Lamperti transform of Y* .

An equivalent definition of S and T , in terms of ξ instead of Y , is given by taking $T(s) = \int_0^s \exp(\alpha \xi_u) du$ and S as its inverse. Then,

$$Y_t = \exp(\xi_{S(t)}) \quad (10)$$

for all $t \geq 0$, and this shows that the Lamperti transform is a bijection.

4.1 The path-censored stable process

Let X be the stable process defined in section 4. In [26], the present authors considered a ‘path-censored’ version of the stable process, formed by erasing the time spent in the negative half-line. To be precise, define

$$A_t = \int_0^t \mathbb{1}_{\{X_s > 0\}} ds, \quad t \geq 0,$$

and let $\gamma(t) = \inf\{s \geq 0 : A_s > t\}$ be its right-continuous inverse. Also define

$$T_0 = \inf\{t \geq 0 : X_{\gamma(t)} = 0\},$$

which is finite or infinite a.s. accordingly as $\alpha > 1$ or $\alpha \leq 1$. Then, the process

$$Y_t = X_{\gamma(t)} \mathbb{1}_{\{t < T_0\}}, \quad t \geq 0,$$

is a pssMp, called the *path-censored stable process*.

In Theorems 5.3 and 5.5 of [26], it is shown that the Laplace exponent ψ^Y of the Lamperti transform ξ^Y associated to Y is given by

$$\psi^Y(z) = \frac{\Gamma(\alpha\rho - z)}{\Gamma(-z)} \frac{\Gamma(1 - \alpha\rho + z)}{\Gamma(1 - \alpha + z)},$$

and there it is remarked that when $\alpha \leq 1$, this process is in the hypergeometric class with parameters

$$(\beta, \gamma, \hat{\beta}, \hat{\gamma}) = (1, \alpha\rho, 1 - \alpha, \alpha\hat{\rho}).$$

It is readily seen from our definition that, when $\alpha > 1$, the process ξ^Y is in the extended hypergeometric class, with the same set of parameters.

From the Lamperti transform we know that

$$I(-\alpha\xi^Y) = \inf\{u \geq 0 : Y_u = 0\} = \int_0^{T_0} \mathbb{1}_{\{X_t > 0\}} dt,$$

where the latter is the occupation time of $(0, \infty)$ up to first hitting zero for the stable process. This motivates the following proposition, whose proof is a direct application of Proposition 4.

Proposition 5. *The Mellin transform of the random variable $I(-\alpha\xi^Y)$ is given, for $\operatorname{Re} s \in (\rho - 1/\alpha, 2 - 1/\alpha)$, by*

$$\mathcal{M}(s) = c \frac{G(2/\alpha - 1 + s; 1/\alpha)}{G(2/\alpha - \rho + s; 1/\alpha)} \frac{G(1/\alpha + \rho + 1 - s; 1/\alpha)}{G(1/\alpha + 1 - s; 1/\alpha)} \frac{\Gamma(1/\alpha - \rho + s)}{\Gamma(\rho + 1 - s)} \Gamma(2 - 1/\alpha - s),$$

where c is a normalising constant such that $\mathcal{M}(1) = 1$.

Remark 6. When X is in the class $\mathcal{C}_{k,l}$ introduced by Doney [12], which is to say

$$\rho + k = l/\alpha,$$

for $k, l \in \mathbb{Z}$, equivalent expressions in terms of gamma and trigonometric functions may be found via repeated application of certain identities of the double-gamma function; see, for example, [17, equations (19) and (20)].

For example, when $k, l \geq 0$, one has

$$\begin{aligned} \mathcal{M}(s) &= c(-1)^l (2\pi)^{l(1/\alpha-1)} (1/\alpha)^{l(1-2/\alpha)} \frac{\Gamma(\frac{1-l}{\alpha} + k + s)}{\Gamma(\frac{l}{\alpha} + 1 - k - s)} \frac{\Gamma(2 - 1/\alpha - s)}{\Gamma(2 - l - \alpha - \alpha s)} \\ &\quad \times \prod_{j=1}^l \Gamma(j/\alpha + 1 - s) \Gamma(2/\alpha - (j/\alpha + 1 - s)) \prod_{i=0}^{k-1} \frac{\sin(\pi\alpha(s+i))}{\pi}, \end{aligned}$$

and when $k < 0$, $l \geq 0$,

$$\begin{aligned} \mathcal{M}(s) &= c(-1)^l (2\pi)^{l(1/\alpha-1)} (1/\alpha)^{l(1-2/\alpha)} \\ &\quad \times \frac{\Gamma(\frac{1-l}{\alpha} + k + s) \Gamma(2 - 1/\alpha - s) \Gamma(l + 1 + \alpha - \alpha s) \Gamma(2 - l + \alpha k + \alpha s)}{\Gamma(\frac{l}{\alpha} + 1 - k - s)} \\ &\quad \times \prod_{j=1}^l \Gamma(j/\alpha + 1 - s) \Gamma(2/\alpha - (j/\alpha + 1 - s)) \prod_{i=2}^{-k-1} \frac{\pi}{\sin(\pi\alpha(s-i))}. \end{aligned}$$

Similar expressions may be obtained when $k \geq 0$, $l < 0$ and $k, l < 0$.

4.2 The radial part of the symmetric stable process

If X is a symmetric stable process—that is, $\rho = 1/2$ —then the process

$$R_t = |X_t|, \quad t \geq 0,$$

is a pssMp, which we call the *radial part* of X . The Lamperti transform, ξ^R , of this process was studied by Caballero et al. [7] in dimension d ; these authors computed the Wiener–Hopf factorisation of ξ^R under the assumption $\alpha < d$, finding that the process is a hypergeometric Lévy process. Using the extended hypergeometric class, we extend this result, in one dimension, by finding the Wiener–Hopf factorisation when $\alpha > 1$.

In Kuznetsov et al. [21], the following theorem is proved using the work of Caballero et al. [7].

Theorem 7 (Laplace exponent). *The Laplace exponent of the Lévy process $2\xi^R$ is given by*

$$\psi^R(2z) = -2^\alpha \frac{\Gamma(\alpha/2 - z)}{\Gamma(-z)} \frac{\Gamma(1/2 + z)}{\Gamma((1 - \alpha)/2 + z)}. \quad (11)$$

We now identify the Wiener–Hopf factorisation of ξ^R , which will depend on the value of α . However, note the factor 2^α in (11). In the context of the Wiener–Hopf factorisation, we could ignore this factor by picking an appropriate normalisation of local time; however, another approach is as follows.

Let us write $R' = \frac{1}{2}R$, and denote by $\xi^{R'}$ the Lamperti transform of R' . Then the scaling of space on the level of the self-similar process is converted by the Lamperti transform into a scaling of time, so that $\xi_s^R = \log 2 + \xi_{s2^\alpha}^{R'}$. In particular, if we write ψ' for the characteristic exponent of $\xi^{R'}$, it follows that $\psi' = 2^{-\alpha}\psi^R$. This allows us to disregard the inconvenient constant factor in (11), if we work with $\xi^{R'}$ instead of ξ^R .

The following corollary is now simple when we bear in mind the hypergeometric class of Lévy processes introduced in section 2. We emphasise that this Wiener–Hopf factorisation was derived by different methods in [7, Theorem 7] for $\alpha < 1$, though not $\alpha = 1$.

Corollary 8 (Wiener–Hopf factorisation, $\alpha \in (0, 1]$). *The Wiener–Hopf factorisation of $2\xi^{R'}$ when $\alpha \in (0, 1]$ is given by*

$$\psi'(2z) = -\frac{\Gamma(\alpha/2 - z)}{\Gamma(-z)} \times \frac{\Gamma(1/2 + z)}{\Gamma(1 - \alpha)/2 + z)}$$

and $2\xi^{R'}$ is a Lévy process of the hypergeometric class with parameters

$$(\beta, \gamma, \hat{\beta}, \hat{\gamma}) = (1, \alpha/2, (1 - \alpha)/2, \alpha/2).$$

Proof. It suffices to compare the characteristic exponent with that of a hypergeometric Lévy process. \square

When $\alpha > 1$, the process $\xi^{R'}$ is not a hypergeometric Lévy process; however, it is in the extended hypergeometric class, and we therefore have the following result, which is new.

Theorem 9 (Wiener–Hopf factorisation, $\alpha \in (1, 2)$). *The Wiener–Hopf factorisation of $2\xi^{R'}$ when $\alpha \in (1, 2)$ is given by*

$$\psi'(2z) = -\left(\frac{\alpha - 1}{2} - z\right) \frac{\Gamma(\alpha/2 - z)}{\Gamma(1 - z)} \times z \frac{\Gamma(1/2 + z)}{\Gamma((3 - \alpha)/2 + z)} \quad (12)$$

and $2\xi^{R'}$ is a Lévy process in the extended hypergeometric class, with parameters

$$(\beta, \gamma, \hat{\beta}, \hat{\gamma}) = (1, \alpha/2, (1 - \alpha)/2, \alpha/2).$$

Proof. Simply use Theorem 7; using the formula $x\Gamma(x) = \Gamma(x+1)$ yields (12). That this is indeed the Wiener-Hopf factorisation follows once we recognise $2\xi^{R'}$ as a process in the extended hypergeometric class, and apply Proposition 1. \square

As an illustration of the utility of the extended hypergeometric class, we will now derive an expression for the Mellin transform of the exponential functional for the dual process $-\xi^{R'}$. This quantity is linked by the Lamperti representation to the hitting time of zero for X ; see section 4. In particular, if

$$T_0 = \inf\{t \geq 0 : X_t = 0\},$$

we have that

$$T_0 = \int_0^\infty e^{\alpha\xi_t^R} dt = \int_0^\infty e^{\alpha\xi_{2^\alpha t}^{R'}} dt = 2^{-\alpha} \int_0^\infty e^{\alpha\xi_s^{R'}} ds = 2^{-\alpha} I(-\alpha\xi^{R'}). \quad (13)$$

Since $-2\xi^{R'}$ is an extended hypergeometric Lévy process with parameters $(\frac{\alpha+1}{2}, \frac{\alpha}{2}, 0, \frac{\alpha}{2})$, which drifts to $+\infty$, we may apply the theory just developed to compute the Mellin transform of $I(-\alpha\xi^{R'})$. Denote this by \mathcal{M} ; that is,

$$\mathcal{M}(s) = \mathbb{E}[I(-\alpha\xi^{R'})^{s-1}],$$

for some range of $s \in \mathbb{C}$ to be determined.

Proposition 10. *For $\operatorname{Re} s \in (-1/\alpha, 2 - 1/\alpha)$,*

$$\begin{aligned} \mathbb{E}_1[T_0^{s-1}] &= 2^{-\alpha(s-1)} \mathcal{M}(s) \\ &= 2^{-\alpha(s-1)} \frac{\sqrt{\pi}}{\Gamma(\frac{1}{\alpha})\Gamma(1-\frac{1}{\alpha})} \frac{\Gamma(1+\frac{\alpha}{2}-\frac{\alpha s}{2})}{\Gamma(\frac{1-\alpha}{2}+\frac{\alpha s}{2})} \Gamma(\frac{1}{\alpha}-1+s) \frac{\Gamma(2-\frac{1}{\alpha}-s)}{\Gamma(2-s)}. \end{aligned} \quad (14)$$

Proof. Let ζ be a hypergeometric Lévy process with parameters $(\frac{\alpha-1}{2}, \frac{\alpha}{2}, 1, \frac{\alpha}{2})$, and denote by $\widetilde{\mathcal{M}}$ the Mellin transform of the exponential functional $I(\alpha/2 \cdot \zeta)$, which is known to be finite for $\operatorname{Re} s \in (0, 1 + 2/\alpha)$ by the argument in the proof of Proposition 4.

We can then use Proposition 4 to make the following calculation, provided that $\operatorname{Re} s \in (0, 2 - 1/\alpha)$. Here G is the double gamma function, as defined in [17, S§3], and we use [17, equation (25)] in the third line and the identity $x\Gamma(x) = \Gamma(x+1)$ in the final line. For normalisation constants C (and C') to be determined, we have

$$\begin{aligned} \mathcal{M}(s) &= C \widetilde{\mathcal{M}}(s) \frac{\Gamma(\frac{1}{\alpha}+s)}{\Gamma(s)} \frac{\Gamma(2-\frac{1}{\alpha}-s)}{\Gamma(2-s)} \\ &= C \frac{G(\frac{3}{\alpha}-1+s; \frac{2}{\alpha})}{G(\frac{3}{\alpha}+s; \frac{2}{\alpha})} \frac{G(\frac{2}{\alpha}+2-s; \frac{2}{\alpha})}{G(\frac{2}{\alpha}+1-s; \frac{2}{\alpha})} \Gamma(s) \frac{\Gamma(\frac{1}{\alpha}+s)}{\Gamma(s)} \frac{\Gamma(2-\frac{1}{\alpha}-s)}{\Gamma(2-s)} \\ &= C \frac{\Gamma(1+\frac{\alpha}{2}-\frac{\alpha s}{2})}{\Gamma(\frac{3-\alpha}{2}+\frac{\alpha s}{2})} \Gamma(\frac{1}{\alpha}+s) \frac{\Gamma(2-\frac{1}{\alpha}-s)}{\Gamma(2-s)} \\ &= C' \frac{\Gamma(1+\frac{\alpha}{2}-\frac{\alpha s}{2})}{\Gamma(\frac{1-\alpha}{2}+\frac{\alpha s}{2})} \Gamma(\frac{1}{\alpha}-1+s) \frac{\Gamma(2-\frac{1}{\alpha}-s)}{\Gamma(2-s)}. \end{aligned}$$

The condition $\mathcal{M}(1) = 1$ means that we can calculate

$$C' = \frac{\sqrt{\pi}}{\Gamma(\frac{1}{\alpha})\Gamma(1 - \frac{1}{\alpha})}, \quad (15)$$

and this gives the Mellin transform explicitly, for $\operatorname{Re} s \in (0, 2 - 1/\alpha)$.

We now expand the domain of \mathcal{M} . Note that, in contrast to the general case of Proposition 4, the right-hand side of (14) is well-defined when $\operatorname{Re} s \in (-1/\alpha, 2 - 1/\alpha)$, and is indeed analytic in this region. (The reason for this difference is the cancellation of a simple pole and zero at the point 0.) Theorem 2 of [28] shows that, if the Mellin transform of a probability measure is analytic in a neighbourhood of the point $1 \in \mathbb{C}$, then it is analytic in a strip $\operatorname{Re} s \in (a, b)$, where $-\infty \leq a < 1 < b \leq \infty$; and furthermore, the function has singularities at a and b , if they are finite. It then follows that the right-hand side of (14) must actually be equal to \mathcal{M} in all of $\operatorname{Re} s \in (-1/\alpha, 2 - 1/\alpha)$, and this completes the proof. \square

We remark that the distribution of T_0 has been characterised previously by Yano et al. [37] and Cordero [11], using rather different methods; and the Mellin transform above was also obtained, again via the Lamperti transform but without the extended hypergeometric class, in Kuznetsov et al. [21].

It is also fairly straightforward to produce the following hitting distribution. Define

$$\sigma_{-1}^1 = \inf\{t \geq 0 : X_t \notin [-1, 1]\},$$

the *first exit time* of $[-1, 1]$ for X . We give the distribution of the position of the symmetric stable process X at time σ_{-1}^1 , provided this occurs before X hits zero. Note that when $\alpha \in (0, 1]$, the process does not hit zero, so the distribution is simply that found by Rogozin [33].

Proposition 11. *Let X be the symmetric stable process with $\alpha \in (1, 2)$. Then, for $|x| < 1$, $y > 1$,*

$$\begin{aligned} & P_x(|X_{\sigma_{-1}^1}| \in dy; \sigma_{-1}^1 < T_0)/dy \\ &= \frac{\sin(\pi\alpha/2)}{\pi} |x|(1 - |x|)^{\alpha/2} y^{-1} (y - 1)^{-\alpha/2} (y - |x|)^{-1} \\ &+ \frac{1}{2} \frac{\sin(\pi\alpha/2)}{\pi} y^{-1} (y - 1)^{-\alpha/2} |x|^{(\alpha-1)/2} \int_0^{1-|x|} t^{\alpha/2-1} (1 - t)^{-(\alpha-1)/2} dt. \end{aligned}$$

Proof. The starting point of the proof is the ‘second factorisation identity’ [22, Exercise 6.7],

$$\int_0^\infty e^{-qz} \mathbb{E}\left[e^{-\beta(\xi_{S_z^+} - z)}; S_z^+ < \infty\right] dx = \frac{\kappa(q) - \kappa(\beta)}{(q - \beta)\kappa(q)}, \quad q, \beta > 0,$$

where

$$S_z^+ = \inf\{t \geq 0 : \xi_t > z\}.$$

We now invert in q and z , in that order; this is a lengthy but routine calculation, and we omit it. We then apply the Lamperti transform: if $g(z, \cdot)$ is the density of the measure $\mathbb{P}(\xi_{S_z^+} - z \in \cdot; S_z^+ < \infty)$, then

$$P_x(|X_{\sigma_{-1}^1}| \in dy; \sigma_{-1}^1 < T_0) = y^{-1}g(\log|x|^{-1}, \log y),$$

and this completes the proof. \square

The following hitting probability emerges after integrating in the above proposition.

Corollary 12. *For $|x| < 1$,*

$$P_x(T_0 < \sigma_{-1}^1) = (1 - |x|)^{\alpha/2} - \frac{1}{2}|x|^{(\alpha-1)/2} \int_0^{1-|x|} t^{\alpha/2-1} (1-t)^{-(\alpha-1)/2} dt.$$

Finally, it is not difficult to produce the following slightly more general result. Applying the Markov property at time T_0 gives

$$\begin{aligned} P_x(X_{\sigma_{-1}^1} \in dy; \sigma_{-1}^1) &= P_x(X_{\sigma_{-1}^1} \in dy) - P_x(X_{\sigma_{-1}^1} \in dy; T_0 < \sigma_{-1}^1) \\ &= P_x(X_{\sigma_{-1}^1} \in dy) - P_x(T_0 < \sigma_{-1}^1) P_0(X_{\sigma_{-1}^1} \in dy). \end{aligned}$$

The hitting distributions on the right-hand side were found by Rogozin [33], and substituting yields the following corollary.

Corollary 13. *For $|x| < 1$, $|y| > 1$,*

$$\begin{aligned} &P_x(X_{\sigma_{-1}^1} \in dy; \sigma_{-1}^1 < T_0)/dy \\ &= \frac{\sin(\pi\alpha/2)}{\pi} (1-x)^{\alpha/2} (1+x)^{\alpha/2} (y-1)^{-\alpha/2} (y+1)^{-\alpha/2} (y-x)^{-1} \\ &\quad - \left[(1-|x|)^{\alpha/2} - \frac{1}{2}|x|^{(\alpha-1)/2} \int_0^{1-|x|} t^{\alpha/2-1} (1-t)^{-(\alpha-1)/2} dt \right] \\ &\quad \times \frac{\sin(\pi\alpha/2)}{\pi} (y-1)^{-\alpha/2} (y+1)^{-\alpha/2} y^{-1}. \end{aligned}$$

4.3 The radial part of the symmetric stable process conditioned to avoid zero

Above, we computed the Lamperti transform of the pssMp $R' = \frac{1}{2}|X|$, where X was a symmetric stable process, and called it $\xi^{R'}$. In this section we consider instead the symmetric stable process conditioned to avoid zero, and obtain its Lamperti transform.

In [30], Pantí shows (among many other results) that the function

$$h(x) = \begin{cases} -\Gamma(1-\alpha) \frac{\sin(\pi\alpha\hat{\rho})}{\pi} x^{\alpha-1}, & x > 0, \\ -\Gamma(1-\alpha) \frac{\sin(\pi\alpha\rho)}{\pi} x^{\alpha-1}, & x < 0, \end{cases}$$

is invariant for the stable process killed upon hitting zero, and defines the family of measures $(P_x^\dagger)_{x \neq 0}$ given via the Doob h -transform:

$$P_x^\dagger(\Lambda) = \frac{1}{h(x)} E_x[h(X_t) \mathbb{1}_\Lambda; t < T_0], \quad x \neq 0,$$

for $\Lambda \in \mathcal{F}_t = \sigma(X_s, s \leq t)$. In [30] it is also shown that the laws P_x^\dagger arise as limits of the stable process conditioned not to have hit zero up to an exponential time of rate q , as $q \downarrow 0$. The canonical process associated with the laws $(P_x^\dagger)_{x \neq 0}$ is therefore called the *stable process conditioned to avoid zero*, and we shall denote it by X^\dagger .

Consider now the process $R^\dagger = \frac{1}{2}|X^\dagger|$. This is a pssMp, and we may consider its Lamperti transform, which we will denote by ξ^\dagger . The characteristics of the generalised Lamperti representation of X^\dagger have been computed explicitly in [10], and the Laplace exponent, ψ^\dagger , of ξ^\dagger could be computed from this information; however, the harmonic transform gives us the following straightforward relationship between Laplace exponents:

$$\psi^\dagger(z) = \psi'(z + \alpha - 1).$$

This allows us to calculate

$$\psi^\dagger(2z) = -\frac{\Gamma(1/2 - z)}{\Gamma((1 - \alpha)/2 - z)} \frac{\Gamma(\alpha/2 + z)}{\Gamma(z)},$$

which demonstrates that $2\xi^\dagger$ is a process in the extended hypergeometric class, with parameters

$$(\beta, \gamma, \hat{\beta}, \hat{\gamma}) = ((\alpha + 1)/2, \alpha/2, 0, \alpha/2).$$

The present authors and A. Kuznetsov previously computed ψ^\dagger in [21], where we also observed that the process ξ^\dagger is the dual Lévy process to $\xi^{R'}$, and remarked that this implies a certain time-reversal relation between R and R^\dagger ; see [8, §2].

5 Concluding remarks

In this section, we offer some comments on how our approach may be adapted in order to offer new insight on an existing class of processes, the Lamperti-stable processes. These were defined in general in the work of Caballero et al. [6], and the one-dimensional Lamperti-stable processes are defined as follows. We say that a Lévy process ξ is in the *Lamperti-stable class* if it has no Gaussian component and its Lévy measure has density

$$\pi(x) = \begin{cases} \mathbf{c}_+ e^{\beta x} (e^x - 1)^{-(\alpha+1)} dx, & x > 0, \\ \mathbf{c}_- e^{-\delta x} (e^{-x} - 1)^{-(\alpha+1)} dx, & x < 0, \end{cases}$$

for some choice of parameters $(\alpha, \beta, \delta, \mathbf{c}_+, \mathbf{c}_-)$ such that $\alpha \in (0, 2)$ and $\beta, \delta, \mathbf{c}_+, \mathbf{c}_- \geq 0$.

The one-dimensional Lamperti-stable processes form a proper subclass of the β -class of Lévy processes of Kuznetsov [15]. It was observed in [17] that there is an intersection

between the hypergeometric class and the Lamperti-stable class. In particular, the Lamperti representations of killed and conditioned stable processes (see [5]) fall within the hypergeometric class; and generally speaking, setting $\beta = \hat{\beta}$ in the hypergeometric class and choosing $\gamma, \hat{\gamma}$ as desired, one obtains a Lamperti-stable process.

However, not all Lamperti-stable processes may be obtained in this way, and we now outline how the ideas developed in this work can be used to characterise another subset of the Lamperti-stable processes.

Define the set of parameters

$$\mathcal{A}_{\text{EHL}} = \{\beta \in [1, 2], \gamma \in (1, 2), \hat{\gamma} \in (-1, 0)\}$$

and for $(\beta, \gamma, \hat{\gamma}) \in \mathcal{A}_{\text{EHL}}$, let

$$\psi(z) = \frac{\Gamma(1 - \beta + \gamma - z) \Gamma(\beta + \hat{\gamma} + z)}{\Gamma(2 - \beta - z) \Gamma(\beta + z)}.$$

Note that this is the negative of the usual hypergeometric Laplace exponent, with $\beta = \hat{\beta}$. We claim that the following proposition holds.

Proposition 14. *There exists a Lévy process ξ with Laplace exponent ψ . Its Wiener–Hopf factorisation $\psi(z) = -\kappa(-z)\hat{\kappa}(z)$ is given by the components*

$$\kappa(z) = \frac{\Gamma(1 - \beta + \gamma + z)}{\Gamma(2 - \beta + z)}, \quad \hat{\kappa}(z) = (\beta - 1 + z) \frac{\Gamma(\beta + \hat{\gamma} + z)}{\Gamma(\beta + z)}.$$

The ascending ladder height process is a Lamperti-stable subordinator, and the descending factor satisfies

$$\hat{\kappa}(z) = (\mathcal{T}_{\beta-1}v)^*(z), \quad v(z) = \frac{\Gamma(1 + z)}{\Gamma(1 + \hat{\gamma} + z)}.$$

Here, v is the Laplace exponent of a Lamperti-stable subordinator.

The process ξ has no Gaussian component and has a Lévy density given by

$$\pi(x) = \begin{cases} \frac{\Gamma(\gamma + \hat{\gamma} + 1)}{\Gamma(1 + \gamma)\Gamma(-\gamma)} e^{(\beta + \hat{\gamma})x} (e^x - 1)^{-(\gamma + \hat{\gamma} + 1)}, & x > 0, \\ \frac{\Gamma(\gamma + \hat{\gamma} + 1)}{\Gamma(1 + \hat{\gamma})\Gamma(-\hat{\gamma})} e^{-(1 - \beta + \gamma)x} (e^{-x} - 1)^{-(\gamma + \hat{\gamma} + 1)}, & x < 0. \end{cases}$$

Thus, ξ falls within the Lamperti-stable class, and

$$(\alpha, \beta, \delta) = (\gamma + \hat{\gamma}, \beta + \hat{\gamma}, 1 - \beta + \gamma).$$

The proposition may be proved in much the same way as Proposition 1, first using the theory of philanthropy to prove existence, and then the theory of meromorphic Lévy processes to deduce the Lévy measure.

We have thus provided an explicit spatial Wiener–Hopf factorisation of a subclass of Lamperti-stable processes disjoint from that given by the hypergeometric processes.

Acknowledgements We would like to thank the anonymous referee for their thorough reading and helpful comments, which have improved the article.

References

- [1] J. Bertoin. *Lévy processes*, volume 121 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1996. ISBN 0-521-56243-0.
- [2] J. Bertoin and M. Yor. The entrance laws of self-similar Markov processes and exponential functionals of Lévy processes. *Potential Anal.*, 17(4):389–400, 2002. ISSN 0926-2601. doi:10.1023/A:1016377720516.
- [3] J. Bertoin and M. Yor. Exponential functionals of Lévy processes. *Probab. Surv.*, 2:191–212, 2005. ISSN 1549-5787. doi:10.1214/154957805100000122.
- [4] R. M. Blumenthal and R. K. Gettoor. *Markov processes and potential theory*. Pure and Applied Mathematics, Vol. 29. Academic Press, New York, 1968.
- [5] M. E. Caballero and L. Chaumont. Conditioned stable Lévy processes and the Lamperti representation. *J. Appl. Probab.*, 43(4):967–983, 2006. ISSN 0021-9002. doi:10.1239/jap/1165505201.
- [6] M. E. Caballero, J. C. Pardo, and J. L. Pérez. On Lamperti stable processes. *Probab. Math. Statist.*, 30(1):1–28, 2010. ISSN 0208-4147.
- [7] M. E. Caballero, J. C. Pardo, and J. L. Pérez. Explicit identities for Lévy processes associated to symmetric stable processes. *Bernoulli*, 17(1):34–59, 2011. ISSN 1350-7265. doi:10.3150/10-BEJ275.
- [8] L. Chaumont and J. C. Pardo. The lower envelope of positive self-similar Markov processes. *Electron. J. Probab.*, 11:no. 49, 1321–1341, 2006. ISSN 1083-6489. doi:10.1214/EJP.v11-382.
- [9] L. Chaumont, A. E. Kyprianou, and J. C. Pardo. Some explicit identities associated with positive self-similar Markov processes. *Stochastic Process. Appl.*, 119(3):980–1000, 2009. ISSN 0304-4149. doi:10.1016/j.spa.2008.05.001.
- [10] L. Chaumont, H. Pantí, and V. Rivero. The Lamperti representation of real-valued self-similar Markov processes. *Bernoulli*, 19(5B):2494–2523, 2013. ISSN 1350-7265. doi:10.3150/12-BEJ460.
- [11] F. Cordero. *On the excursion theory for the symmetric stable Lévy processes with index $\alpha \in]1, 2]$ and some applications*. PhD thesis, Université Pierre et Marie Curie – Paris VI, 2010.
- [12] R. A. Doney. On Wiener-Hopf factorisation and the distribution of extrema for certain stable processes. *Ann. Probab.*, 15(4):1352–1362, 1987. ISSN 0091-1798.

- [13] A. V. Gnedin. Regeneration in random combinatorial structures. *Probab. Surv.*, 7: 105–156, 2010. ISSN 1549-5787. doi:10.1214/10-PS163.
- [14] I. S. Gradshteyn and I. M. Ryzhik. *Table of integrals, series, and products*. Elsevier/Academic Press, Amsterdam, seventh edition, 2007. ISBN 978-0-12-373637-6. Translated from the Russian. Translation edited and with a preface by Alan Jeffrey and Daniel Zwillinger.
- [15] A. Kuznetsov. Wiener-Hopf factorization and distribution of extrema for a family of Lévy processes. *Ann. Appl. Probab.*, 20(5):1801–1830, 2010. ISSN 1050-5164. doi:10.1214/09-AAP673.
- [16] A. Kuznetsov. Wiener-Hopf factorization for a family of Lévy processes related to theta functions. *J. Appl. Probab.*, 47(4):1023–1033, 2010. ISSN 0021-9002.
- [17] A. Kuznetsov and J. C. Pardo. Fluctuations of stable processes and exponential functionals of hypergeometric Lévy processes. *Acta Appl. Math.*, 123:113–139, 2013. ISSN 0167-8019. doi:10.1007/s10440-012-9718-y.
- [18] A. Kuznetsov, A. E. Kyprianou, J. C. Pardo, and K. van Schaik. A Wiener-Hopf Monte Carlo simulation technique for Lévy processes. *Ann. Appl. Probab.*, 21(6): 2171–2190, 2011. ISSN 1050-5164. doi:10.1214/10-AAP746.
- [19] A. Kuznetsov, A. E. Kyprianou, and J. C. Pardo. Meromorphic Lévy processes and their fluctuation identities. *Ann. Appl. Probab.*, 22(3):1101–1135, 2012. doi:10.1214/11-AAP787.
- [20] A. Kuznetsov, A. E. Kyprianou, and V. Rivero. The theory of scale functions for spectrally negative Lévy processes. In *Lévy Matters II*, Lecture Notes in Mathematics, pages 97–186. Springer, Berlin and Heidelberg, 2013. ISBN 978-3-642-31406-3. doi:10.1007/978-3-642-31407-0_2.
- [21] A. Kuznetsov, A. E. Kyprianou, J. C. Pardo, and A. R. Watson. The hitting time of zero for a stable process. *Electron. J. Probab.*, 19:no. 30, 1–26, 2014. ISSN 1083-6489. doi:10.1214/EJP.v19-2647.
- [22] A. E. Kyprianou. *Introductory lectures on fluctuations of Lévy processes with applications*. Universitext. Springer-Verlag, Berlin, 2006. ISBN 978-3-540-31342-7.
- [23] A. E. Kyprianou and P. Patie. A Ciesielski-Taylor type identity for positive self-similar Markov processes. *Ann. Inst. Henri Poincaré Probab. Stat.*, 47(3):917–928, 2011. ISSN 0246-0203. doi:10.1214/10-AIHP398.
- [24] A. E. Kyprianou and V. Rivero. Special, conjugate and complete scale functions for spectrally negative Lévy processes. *Electron. J. Probab.*, 13:1672–1701, 2008. ISSN 1083-6489. doi:10.1214/EJP.v13-567.

- [25] A. E. Kyprianou, J. C. Pardo, and V. Rivero. Exact and asymptotic n -tuple laws at first and last passage. *Ann. Appl. Probab.*, 20(2):522–564, 2010. ISSN 1050-5164. doi:10.1214/09-AAP626.
- [26] A. E. Kyprianou, J. C. Pardo, and A. R. Watson. Hitting distributions of α -stable processes via path censoring and self-similarity. *Ann. Probab.*, 42(1):398–430, 2014. ISSN 0091-1798. doi:10.1214/12-AOP790.
- [27] J. Lamperti. Semi-stable Markov processes. I. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete*, 22:205–225, 1972.
- [28] E. Lukacs and O. Szász. On analytic characteristic functions. *Pacific J. Math.*, 2: 615–625, 1952. ISSN 0030-8730.
- [29] F. W. J. Olver. *Asymptotics and special functions*. Academic Press, New York–London, 1974. Computer Science and Applied Mathematics.
- [30] H. Pantí. On Lévy processes conditioned to avoid zero. arXiv:1304.3191v1 [math.PR], 2012.
- [31] J. C. Pardo. The upper envelope of positive self-similar Markov processes. *J. Theoret. Probab.*, 22(2):514–542, 2009. ISSN 0894-9840. doi:10.1007/s10959-008-0152-z.
- [32] V. Rivero. Recurrent extensions of self-similar Markov processes and Cramér’s condition. II. *Bernoulli*, 13(4):1053–1070, 2007. ISSN 1350-7265. doi:10.3150/07-BEJ6082.
- [33] B. A. Rogozin. The distribution of the first hit for stable and asymptotically stable walks on an interval. *Theory of Probability and its Applications*, 17(2):332–338, 1972. ISSN 0040585X. doi:10.1137/1117035.
- [34] R. Song and Z. Vondraček. Potential theory of special subordinators and subordinate killed stable processes. *J. Theoret. Probab.*, 19(4):817–847, 2006. ISSN 0894-9840. doi:10.1007/s10959-006-0045-y.
- [35] V. Vigon. *Simplifiez vos Lévy en titillant la factorisation de Wiener–Hopf*. PhD thesis, INSA de Rouen, 2002.
- [36] J. Vuolle-Apiala. Itô excursion theory for self-similar Markov processes. *Ann. Probab.*, 22(2):546–565, 1994. ISSN 0091-1798.
- [37] K. Yano, Y. Yano, and M. Yor. On the laws of first hitting times of points for one-dimensional symmetric stable Lévy processes. In *Séminaire de Probabilités XLII*, volume 1979 of *Lecture Notes in Math.*, pages 187–227. Springer, Berlin, 2009. doi:10.1007/978-3-642-01763-6_8.